The background of the slide is a high-angle aerial photograph of a coastal region. The left side is dominated by a deep blue ocean with visible white-capped waves. The right side shows a green landscape with rolling hills, some brown patches, and a winding road or path. The overall scene is a mix of natural coastal beauty and human-made infrastructure.

Description of Recent Updates to the NRL Coupled Ocean Data Assimilation System (NCODA)

*James Cummings
Oceanography Division
Naval Research Laboratory
Monterey, CA*

*HYCOM NOPP GODAE Meeting
RSMAS October 2004*

Report Documentation Page

*Form Approved
OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE OCT 2004	2. REPORT TYPE	3. DATES COVERED 00-00-2004 to 00-00-2004		
4. TITLE AND SUBTITLE Description of Recent Updates to the NRL Coupled Ocean Data Assimilation System (NCODA)			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Oceanography Division, Monterey, CA, 93943			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited				
13. SUPPLEMENTARY NOTES				
14. ABSTRACT				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified		
			18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON



NRL Coupled Ocean Data Assimilation System

Advanced DA Technique Based on Optimal Estimation Theory


- oceanographic version of MVOI method used in NWP systems
Lorenc (1981), Daley (1991)
- simultaneous analysis of five ocean variables: temperature, salinity, geopotential, and u-v velocity components (T , S , Φ , u , v)
- developed as part of the ONR sponsored coupled modeling projects at NRL MRY (COAMPS/NCOM and NOGAPS/POP)

Flexible System

- supports variety of map projections
- performs multi-scale analyses on nested, successively higher resolution grids
- initialize/update ocean forecast model or run stand-alone
 - 2D analyses of sea ice and SST (NWP boundary conditions)
 - 3D temperature and salinity analysis (geostrophic currents)
 - 3D MVOI sequential incremental update cycle

NRL Coupled Ocean Data Assimilation

Sequential Incremental Update Cycle
Analysis-Forecast-Ana



Forecast fields and prediction errors
can be used in the QC of new data



FNMOC Quality Controlled Operational Ocean Observation Data Sources

- **AVHRR GAC Satellite SST** ~800,000 obs/day
 - 8-km resolution (NOAA 16,17 day, night, relaxed day retrievals)
- **GOES Satellite SST** ~3,000,000 obs/day
 - 12-km resolution (GOES 10 day, night retrievals)
- ***In Situ* SST/SSS** ~15,000 obs/day
 - surface ship, fixed and drifting buoys, CMAN, TRACKOB
- **Subsurface Temperature and Salinity Profiles** ~1000 profiles/day
 - XBTs, CTDs (TESACS), Argo floats
 - fixed buoys (TAO, PIRATA), thermistor chain drifting buoys
- **Sea Surface Height Anomaly (SSHA)** ~100,000 obs/day
 - altimeter (GFO, ENVISAT, Jason-1), XBTs, CTDs, Argo Floats
- **Sea Ice Concentration** ~1,000,000 obs/day
 - SSM/I (DMSP F13, F14, F15)

All QC data files available on Monterey GODAE server in real-time



New Ocean Observation Data Sources

- **AVHRR LAC Satellite SST** ~3,200,000 obs/day
 - 2-km resolution (NOAA 17 day, night retrievals)
- **GOES Satellite SST** ~3,000,000 obs/day
 - 12-km resolution (GOES 12 day, night retrievals)
- **AMSR-E Microwave SST** ~180,000 obs/day
 - 25-km resolution (day, night retrievals)
- **AATSR Skin SST** ~90,000 retrievals/day
 - 16-km resolution (day, night retrievals)

AVHRR LAC and GOES 12 satellite SST retrievals operationally available at NAVOCEANO (and GODAE server)

AMSR-E and AATSR satellite SST available on GODAE server



New Analysis Capabilities

- **Irregular Grid**
 - supports global stitched grid (Global NCOM)
 - analysis currently supports Mercator, Polar Stereographic, Lambert Conformal, Spherical, Cartesian grid projections
- **Enhanced QC Tests**
 - SSM/I land contamination
 - flags spurious positive sea ice concentration retrievals near land during summer melt season
 - satellite SST diurnal warming
 - detects warm biased daytime satellite SST retrievals due to diurnal warming events (collocates NWP winds and solar radiation)
 - Argo salinity profile bias correction
 - corrects salinity profile to GDEM 3.0 climate salinity offset at depth
 - profile cross validation
 - checks new profiles with analysis performed using nearby profiles



New Analysis Capabilities

- **Reduced Resolution Assimilation Grid**
 - compute innovations from full model resolution grid; perform analysis on coarser resolution analysis grid
 - cost of assimilation mainly in post multiplication (observation to grid space); reduced resolution analysis grid improves throughput
- **Improved Memory Management**
 - required for large model grids on IBM architecture
- **Wave Model Data Assimilation**
 - altimeter/buoy significant wave height (SWH) data in Wavewatch III
 - FNMOC OPTEST planned for 2004/2005 northern hemisphere winter
 - wave model spectra update using choice of methods (BMRC or NCEP)
 - includes new SWH observation QC module



NCODA Operational Status

- **FNMOC**
 - Ocean QC system operational March 2004
 - NCODA 3D analysis-only capability operational August 2004
 - near global analysis running daily in real-time (27-km mid-latitude resolution Mercator grid)
 - 2D sea ice and SST analyses running daily in real-time on 27-km northern and southern hemisphere polar stereographic grids
- **NAVOCEANO**
 - RTP project transitioned QC and analysis to NAVOCEANO
 - Ocean QC nearing operational status
 - biggest issue is connecting to NAVO data streams
 - global sea ice and SST analyses run daily on NAVOCEANO IBMs to support QC
 - NCODA analysis being used in exercise support within NAVOCEANO




NCODA Plans

- **Conversion to 3DVar - adapt NAVDAS solver to the ocean**
 - allows for greater flexibility for assimilating different observation data types (possibly non-linearly related to the forecast model state)
 - eliminates the need to split the analysis domain into sub-domains (all observations can influence the analysis at every model grid point)
 - provides a clear development path towards more advanced 4D assimilation techniques
- **Velocity Observation Data Assimilation**
 - quality control and pre-processing module development
 - model velocity background checks
 - speed, direction conversion to u,v vectors; analysis grid rotation
 - observation error - instrumentation and representation
 - develop sources of velocity observations in real-time
 - Argo float drift, HF radar, current meters, drifting buoys, wide swath altimetry

HYCOM Tasks

Naval Research Laboratory, Marine Meteorology Division (NRL MMD)


HYCOM



On-Going Research

- **Restructure COAMPS™ CM and Code Structure**
 - ESMF, WRF compliance (FY04-FY06)
 - Adapt Flux Coupler to force HYCOM (FY05+)
 - Add HYCOM as ocean model (FY06+)
- **Test HYCOM fields as IC and LBC for NCOM (FY05)**

NCOM



Needs:

- Daily real-time HYCOM forecast runs to 72 h, w/3 h interval (T, S, u, v, SSH)
- Standardized I/O (filenames, format)
- Areas of interest: Med/Atlantic, east and west Pacific
- Interpolation code (HYCOM-to-NCOM vertical grids)
- HYCOM ported to ESMF

ESMF: **Earth System Modeling Framework**

WRF: **Weather Research and Forecast (Infrastructure)**

COAMPS™ is a trademark of the Naval Research Laboratory